

COSMOLOGICAL PARAMETERS: do we already know the final answer ?

MICHAEL ROWAN-ROBINSON

*Astrophysics group, Blackett Laboratory, Imperial College,
Prince Consort Rd, London SW7 2BZ
E-mail: mrr@ic.ac.uk*

Some of the arguments which support the strong consensus for an $\Omega_o = 0.3$, $\lambda_o = 0.7$ model are reexamined. Corrections for Malmquist bias, local flow and metallicity suggest a revised value for H_o of 63 ± 6 km/s/Mpc, improving the age problems for an $\Omega_o = 1$ universe. The latest CMB results may require a high baryon density and hence new physics, for example a strong lepton asymmetry. Difficulties for the $\Omega_o = 1$ model with cluster evolution, the baryon content of clusters, and the evidence from Type Ia supernovae favouring low Ω_o , $\Lambda > 0$ models, are discussed critically.

1 Introduction

A strong consensus has developed in support of the cosmological parameter set:

$$\Omega_o = 0.3, \lambda_o = 0.7, H_o = 75, t_o = 12.6 \text{ Gyr.}$$

Is this justified ? How certain is the evidence that $\Lambda > 0$? In this paper I test how strong some of the evidence is, and show that the latest CMB results may point towards some interesting radical alternatives.

Some of the problems for an $\Omega_o = 1, \Lambda = 0$ model are:

- observed $H_o t_o$ too high
- Type Ia supernovae imply $\lambda_o > 0$ and CMB gives $\Omega_o + \lambda_o = 1$
- cluster abundance evolution favours low Ω_o
- baryon fraction in clusters favours low Ω_o
- $P(k)$ for galaxies requires $\Gamma \simeq \Omega_o h \simeq 0.2$

However these problems may not be as insuperable as they appear.

2 Hubble constant, H_o

New distances from the HST Key program on Cepheids, and their application to Type Ia and II supernovae, and the Tully-Fisher, $D_n - \sigma$, and surface brightness fluctuation methods, have been reviewed by Freedman (2000) and

Mould et al (2000, and references therein). In these papers the approach is to determine the Hubble constant via different distance methods and then combine these estimates statistically.

The philosophy of the present work is that of Rowan-Robinson (1985 (RR85), 1988), to form weighted mean distances of groups and clusters and then restrict the analysis to high weight distances in order to minimise Malmquist bias

An important ingredient in solving for the Hubble constant is the flow model. Estimation of peculiar velocities is one of the largest remaining uncertainties in H_o . Mould et al (2000) use a simple 3-attractor model. The PSCz flow model of Rowan-Robinson et al (2000), based on 88 groups, 854 clusters and 163 voids, gives much better resolution of the local flow.

For Type Ia supernovae it is important to correct for internal extinction in the host galaxy. Phillips et al (1999) have carried out an important analysis of internal extinction for Type Ia supernovae and I have used these estimates where available, otherwise the RR85 prescription. I use the calibration of Gibson et al (2000), based on 8 Type Ia supernovae with HST Cepheid distances, for which the absolute magnitude at maximum light is then $\langle M_B \rangle = -19.47$.

An important correction to the HST Cepheid distances is for the differences in metallicity between Cepheids in the LMC and the HST program galaxies. Mould et al (2000) estimate that this increases the HST Cepheid distance scale by 4%.

The HST Key program is based on $\mu_o(LMC) = 18.5$, but Feast (2000) estimates that this may be too low by 7%, based on Hipparcos parallaxes (after correction for metallicity effects in Cepheids).

Figure 1 shows how the mean Hubble constant, calculated for groups and clusters with well-determined mean distances, varies with the assumed minimum weight. Inclusion of lower weight cluster and group distances leads to an increased mean Hubble constant, almost certainly the effect of Malmquist bias. The results are shown for the Mould et al (2000) 3-attractor flow model, and for the flow model derived from the PSCz analysis (Rowan-Robinson et al (2000)). The difference between the flow models is not large, but it is significant.

Figure 2 shows the Hubble diagram for groups and clusters with distances less than 400 Mpc. The mean Hubble constant for clusters with weight ($= \Sigma W / \sigma^2$) > 15 is found to be $H_o = 63 \pm 6$ km/s/Mpc (Rowan-Robinson, 2000a, in preparation), after correction for the effect of metallicity (but with $\mu_o(LMC) = 18.5$). I have adopted the uncertainty quoted by Mould et al (2000).




Figure 1: Effect of different minimum weights on mean value of Hubble constant. Filled circles: PSCz flow model (Rowan-Robinson et al 2000). Open circles: 3-attractor model (Mould et al 2000). No correction for metallicity has been applied. The error bars include only statistical uncertainties.

2.1 Comparison with other estimates

Cepheids: Freedman (2000) have summarized the results of the HST Key Program using Cepheids in galaxies out to 20 Mpc to determine the Hubble constant and found $H_o = 74 \pm 8$ km/s/Mpc. Mould et al (2000), with their 3-attractor model for the local flow, find $H_o = 71 \pm 6$ km/s/Mpc. The differences between my estimate and Mould et al's can be assigned as follows: improved flow model $\Delta H_o = -2$, correction for Malmquist bias -2, correction for metallicity -3.

Supernovae Type Ia: Branch (1998) has given an excellent review of the current situation. Initially widely disagreeing estimates of how the absolute magnitude at maximum light depends on decay rate have to some extent converged. Hamuy et al (1996) analyzed a large sample of nearby supernovae and find $H_o = 63 \pm 4.5$ km/s/Mpc. Saha et al (1997) analyzed 7 Type Ia supernovae with HST Cepheid distances and found $H_o = 58 \pm 8$ km/s/Mpc (61 if they included a relationship between M_B and ΔM_{15}). Branch et al (1996) found $H_o = 57 \pm 4$ km/s/Mpc from a colour matched sample of supernovae and Tripp (1997) found $H_o = 60 \pm 5$ km/s/Mpc from a sample matched according to Δm_{15} . Riess et al (1996) used a template method in which multicolour photometry is used to characterize supernovae in a 1-parameter sequence ('MLCS')




Figure 2: Hubble diagram for groups and clusters with weight > 3 .

to obtain $H_o = 64 \pm 6$ km/s/Mpc.

Several groups have brought theoretical models to bear on the determination of the Hubble constant using Type Ia supernovae. Hoflich and Khokhlov (1996) compared 26 supernovae with model light curves and found $H_o = 67 \pm 9$ km/s/Mpc. Branch (1998) suggests this should be revised to 56 ± 5 . The same two authors found $H_o = 55$ if they included a theoretical version of the $M_B - \Delta M_{15}$ relation. Nugent et al (1995) fitted non-LTE model spectra to observations and found $H_o = 60 + 14, -11$. In a recent analysis Tripp and Branch (1999) conclude that the best estimate for the Hubble constant from Type Ia supernovae was $H_o = 62 \pm 5$ km/s/Mpc.

Gravitational lens time delay:

An analysis by Falco et al (1997) of the gravitational lens time delay system 0957+561 gave $H_o = 62 \pm 7$ km/s/Mpc. Koopmans and Fassnacht (1999) use 5 gravitational lenses to determine $H_o = 64 \pm 11$ km/s/Mpc

Sunyaev-Zeldovich effect: Recent work on SZ clusters includes Myers et al (1997), Birkinshaw (1999) and Reese et al (2000). Birkinshaw et al (2000) has given $H_o = 54 \pm 8 \pm 10$ km/s/Mpc for 9 clusters.

To summarize, my estimate for H_o , 63 ± 6 km/s/Mpc agrees well with

independent estimates.

With $\Omega_o = 1, \lambda_o = 0$, the age of the universe becomes 10.2 ± 1.0 Gyr, while for $\lambda_o = 0.7, \Omega_o = 0.3$, the age becomes 15.0 ± 1.4 Gyr. These estimates can be compared with the estimate of Chaboyer et al (1998) for the age of the globular clusters, 11.5 ± 1.3 Gyr, to which must be added the time to form the globular clusters, 0.2-2 Gyr, depending on the redshift of formation. Neither cosmological model would appear to be decisively favoured (or ruled out) by the age estimate.

3 CMB and $\Omega_b h^2$

The pre-Boomerang and Maxima consensus was that $\Omega_o = 0.3, \lambda_o = 0.7, \Omega_b h^2 = 0.019 \pm 0.002, h = 0.68, n = 1$. However this provides a very poor fit to the combined Boomerang and Maxima CMB data. In particular the second peak in the angular power spectrum is found to be too weak for this parameter combination.

The consensus fix (Jaffe et al 2000) is to take a higher baryon density, $\Omega_b h^2 = 0.03$. Unless there is a problem with Big-Bang nucleosynthesis calculations, this would require the deuterium abundance to be significantly lower than even the low estimates of Burles and Tytler (1998a,b) (but see Tegmark et al 2000). It will be interesting to see whether the weakness of the second peak is confirmed in future observations.

This approach also provides no physical explanation of why the vacuum energy density drops at the end of inflation by a factor 10^{110} , only to achieve a significant dynamical role again at the present epoch. It is disappointing that quintessence models, which attempt to give some physical meaning to the evolution of Λ do not seem to fit the data as well as a constant Λ does.

4 A radical alternative: lepton asymmetry

A radical alternative is to consider a strong lepton asymmetry (Harvey and Kolb 1981, Kang and Steigman 1992, Lesgourgues and Pastor 1999, Kinny and Riotto 1999, White et al 2000, Lesgourgues and Peloso 2000, Esposito et al 2000). To maintain charge constancy, this has to be an asymmetry in the neutrino sector. The nucleosynthesis implications have been explored by Kang and Steigman (1992). The enhanced relativistic energy-density speeds up the expansion. Neutrons and protons decouple earlier, which enhances the abundance of neutrons, but this can be compensated by increased neutron decay due to an enhanced ν_e abundance. To get consistency with the observed primordial light element abundances, we then need a higher baryonic density. Lepton asymmetry also has the consequence (similar to the effects

of a positive Λ or a hot dark-matter component) of postponing the epoch of matter-radiation equality, thereby reducing the parameter Γ and improving the fit to the observed $P(k)$ relative to a pure CDM scenario.

The lepton asymmetry can be characterized as an effective number of neutrinos, N_{eff} . The ratio of total energy density in extreme relativistic particles to the energy density in photons, at the epoch of nucleosynthesis is then $\rho_{ER}/\rho_\gamma = 11/4 + 7/8 N_{eff}$. The expansion speed-up factor after $e^\pm$ annihilation can be written $S_o = [1 + 0.135 (N_{eff} - 3)]^{1/2}$. Lesgourgues and Peloso (2000) find a reasonable fit to the Boomerang and Maxima data with $N_{eff} = 6, \Omega_o = 1, \lambda_o = 0, \Omega_b h^2 = 0.028, h = 0.70, n = 1$ and τ (reionization optical depth) = 0.05. Esposito et al (2000) find that the best fit to the Boomerang and Maxima data, retaining consistency with observed light element abundances, would require $N_\nu = 9 \pm 4(2 - \sigma)$.

However to get a good fit to large-scale structure data, eg $P(k)$, we need

$$\Gamma = \Omega_o h / S_o^2 = \Omega_o h / [1 + 0.135(N_\nu - 3)] \simeq 0.2$$

so for $\Omega_o = 1, h = 0.63$ we need $N_\nu \simeq 19$.

The analysis of Esposito et al (2000) suggest that the best compromise would be $N_\nu = 13$, which would give $\Gamma = 0.26$. This would give consistency at the $2 - \sigma$ level with the CMB data, light element abundances and large-scale structure.

A completely different radical alternative consistent with $\Omega_o = 1, \lambda_o = 0, \Omega_b h^2 = 0.019$, and consistent with both the CMB data and large-scale structure data is the proposal for a phase transition during inflation by Barriga et al (2000).

5 The evidence for positive Λ from Type Ia supernovae

There has been a great deal of excitement about the possibility that Type Ia supernovae yield positive evidence that $\Lambda > 0$. Recently two groups have published results on the Hubble diagram for Type Ia supernovae, with over 100 supernovae now discovered at $z > 0.3$ (Schmidt et al 1998, Garnavich et al 1998, Riess et al 1998, Perlmutter et al 1999). Both groups claim that models with positive cosmological constant are preferred, and that models with $\lambda = 0.7, \Omega_o = 0.3$ provide the best fit to the data. The strength of the signal is that Type Ia supernovae at redshift 0.3-0.9 are about 0.25 magnitudes fainter than local supernovae, if an $\Omega_o = 1$ Einstein-de Sitter universe is assumed. Claims that this is a 7-8 σ effect therefore depend on a very precise homogeneity of Type Ia supernovae. The key element in reducing the scatter in Type Ia supernova absolute magnitudes at maximum light has been the correlation between absolute magnitude and decline rate ($M_B - \Delta m_{15}$), discussed above. If




Figure 3: Hubble diagram for SZ clusters (open circles, data from Myers et al 1997, Birkinshaw 1999, Reese et al 2000), Type Ia supernovae (filled circles, data from Perlmutter et al 1998) and gravitational lenses (crosses, data from Koopmans and Fassnacht 1999).

one looks at the paper by Hamuy et al (1996) where this relation is established for 29 local supernovae, one finds that the situation is not quite as impressive as has been presented. It would seem reasonable that to talk about a relationship between the absolute magnitude at maximum and the decline rate over the next 15 days, it would be necessary to have detected the calibrating supernovae prior to maximum. In fact only 10 of the local supernovae were first observed at least one day before maximum. For these 10 there is indeed a $M_B - \Delta m_{15}$ relation, but its significance is much reduced. If we derive the calibration from these 10 local supernovae and apply it to the distant supernovae, the significance of the signal is reduced from the claimed 7-8 σ to only 2-3 σ , depending on which calibration is used (Rowan-Robinson 2000b, in preparation). It appears that the calibrating relation needs to be placed on a much stronger basis with nearby supernovae before it can be used to establish the reality of a cosmological constant. A good test of homogeneity would be to find several supernova in a high redshift cluster.

Fig 3 shows the Hubble diagram, velocity versus luminosity distance, for snIa, SZ clusters and gravitational lens time delay systems. The theoretical curves are shown for an assumed Hubble constant of 63 km/s/Mpc. The evidence from snIa supporting $\Lambda > 0$ comes from supernovae with distances > 3000 Mpc (ie the fainter ones). The gravitational lenses, reaching to significantly greater redshifts, do not support $\Lambda > 0$.

There are also some theoretical uncertainties. Since we do not know for certain whether nearby supernovae are due to white dwarf deflagration or to white dwarf mergers, there is the possibility that the proportion of these two types changes with epoch and this could affect the mean absolute magnitude. Hoflich et al (1998, 2000) also point out that uncertainties and evolution of the initial composition in supernovae can have a significant effect on the determination of cosmological parameters using supernovae.

6 Other problems: the cluster baryon fraction, cluster abundances

- **cluster evolution:** strong negative evolution would be expected in the space density of rich clusters in an $\Omega_o = 1$ universe because growth of structure continues to the present day. This has been claimed not to have been seen by Bahcall et al (1997), Fan et al (1997), Carlberg et al (1997), Eke et al (1998), Bahcall et al (2000). Bahcall et al (2000) estimate that $\Omega_o = 0.16 \pm 0.05$ and argue that $\Omega_o = 1$ is strongly ruled out. However using an X-ray selected sample of clusters, Blanchard et al (2000) estimate $\Omega_o = 0.74 \pm 0.18$, when account is taken of the cluster temperature distribution function.

- **the cluster baryon fraction:** Many groups have confirmed the original finding of White et al (1993) that the baryon content of clusters is too high for an $\Omega_o = 1$ universe (White and Fabian (1995), Loewenstein and Mushotzky (1996), Mulchaey et al (1996), Evrard (1997)), if the baryon density is taken to be $\Omega_b h^2 = 0.019 \pm 0.002$. If however the higher baryonic density implied by the weak second CMB power spectrum peak seen by Boomerang and Maxima is correct, then the problem is not quite so acute. For example Birkinshaw (2000) estimates the $\Omega_b/\Omega_o = 0.07$, and if this is combined with $\Omega_b h^2 = 0.028, h = 0.63$, we deduce that $\Omega_o = 1.008$!

In conclusion there are two viable scenarios consistent with the evidence on cosmological parameters: an $\Omega_o = 1, \lambda = 0$ universe or a $\Omega_o = 0.3, \lambda = 0.7$ universe. Although a number of lines of evidence, including high redshift Type Ia supernovae, favour the latter, the evidence from CMB fluctuations may require new physics to resolve.

References

1. Bahcall N.A., Fan X., Cen R., 1997, ApJ 485, L53
2. Bahcall N.A., Cen R., Dave R., Ostriker J.R., Yu Q., 2000, astro-ph/0002310
3. Barriga J., Gaztanaga E., Santos M.G., Sarkar S., 2000, astro-ph/0011398
4. Birkinshaw M., 1999, Phys. Reports 310, 97
5. Birkinshaw M., 2000, IAU Symposium 201, 'New Cosmological Data and the Values of the FundamentalParameters', eds A.N.Lasenby et al
6. Blanchard A., Sadat R., Bartlett J.G., Le Dour M., 2000, astro-ph/9908037
7. Branch D., 1998, ARAA 36, 17
8. Branch D., Romanishin W., Baron E., 1996, ApJ 465, 73
9. Burles S. and Tytler D., 1998a, ApJ 499, 600; 1998b, ApJ 507, 732
10. Carlberg R.G., Morris S.L., Yee H.K.C., Ellingson E., 1997, ApJ 479, 82
11. Eke V.R., Cole S., Frenk C.S., Henry J.P., 1998, MN 298, 1145
12. Evrard A.E., 1997, MNRAS 292, 289
13. Falco E.E., et al, 1997, ApJ 484, 70
14. Fan X., Bahcall N.A., Cen R., 1997, ApJ 490, L123
15. Feast M., 2000, IAU Symposium 201, 'Cosmological Parameters', 'New Cosmological Data and the Values of the FundamentalParameters', eds A.N.Lasenby et al, astro-ph/0010590
16. Freedman W.L., 2000, IAU Symposium 201, 'Cosmological Parameters'

17. Garnavich P.M. et al, 1998, ApJ 493, L53
18. Gibson B.K. et al, 2000, ApJ, astro-ph/9908149
19. Hamuy M. et al, 1996, AJ 112, 2391
20. Harvey and Kolb, 1981, Phys Rev D24, 2090
21. Hofflich P. and Khokhlov A., 1996, ApJ 457, 500
22. Hofflich P., Wheeler J.C., Thielemann F.K., 1998, ApJ 495, 617
23. Hofflich P., Nomoto K., Umeda H., Wheeler J.C., 2000, astro-ph/9908226
24. Jaffe A.H. et al, 2000, astro-ph/0007333
25. Kang H.-S. and Steigman G., 1992, Nucl.Phys.B 372, 494
26. Kinny W.H. and Riotto A., 1999, Phys.Rev.L 83, 3366
27. Koopmans L.V.E. and Fassnacht C.D., 1999, ApJ 527, 513
28. Lesgourgues J. and Pastor S., 1999, Phys.Rev. D60, 3521
29. Lesgourgues J. and Peloso M., 2000, astro-ph/000412
30. Loewenstein M. and Mushotzky R.L., 1996, ApJ 471, L83
31. Mould J. et al, 2000, astro-ph/9909260
32. Mulchaey J.S. et al, 1996, ApJ 456, 80
33. Myers S.T., Baker J.E., Readhead A.C.S., Herbig T., 1997, ApJ 485, 1
34. Nugent P. et al, 1996, ApJ 485, 812
35. Perlmutter S. et al, 1999, ApJ 517, 565
36. Phillips M.M. et al, 1999, AJ 118, 1766
37. Reese E.D. et al, 2000, ApJ 533, 38
38. Riess A.G., Press W.H., Kirshner R.P., 1996, ApJ 473, 588
39. Riess A.G. et al, 1998, AJ 116, 1009
40. Rowan-Robinson M., 1985, 'The Cosmological Distance Ladder'
(W.H.Freeman)
41. Rowan-Robinson M., 1988, Space Physics Reviews, 48,
42. Rowan-Robinson M. et al, 2000, MN 314, 375
43. Saha A. et al, 1997, ApJ 486, 1
44. Schmidt B.P. et al, 1988, ApJ 507, 46
45. Tegmark M., Zaldarriaga M., Hamilton A.J.S., 2000, astro-ph/0008167
46. Tripp R., 1997, AA 325, 871
47. Tropp R. and Branch D., 1999, ApJ 525, 209
48. White, S.D.M., Navaro J., Evrard A., Frenk C.S., 1993, Nature 366, 429
49. White S.D.M. and Fabian A., 1995, MNRAS 273, 72
50. White M., Scott D. and Pierpaoli E., 2000, astro-ph/0004385